

Anti-c-Kit(CD117) hIgG1 Antibody(briquilimab)

Product information

GM-86543AB-10	10 µg
GM-86543AB-100	100 µg
GM-86543AB-1000	1 mg


Antibody Information

Species Reactivity	Human; Cynomolgus
Clone	briquilimab
Source/Isotype	Monoclonal human IgG1, κ
Application	Flow cytometry; Bioactivity-ELISA
Specificity	Detects c-Kit(CD117)
Gene	c-Kit(CD117)
Other Names	C-Kit, CD117, MASTC, PBT, SCFR
Gene ID	3815 (human); 102137242 (Cynomolgus)
Background	c-Kit, also known as CD117, is an important cell surface protein expressed in stem cells and various types of mature cells in humans and other animals. Encoded by the c-Kit gene, the c-Kit protein plays a role in cell proliferation, differentiation, and survival by binding with the cytokine SCF (stem cell factor). Mutations and abnormal expressions of the c-Kit gene have been associated with various diseases including tumors, cardiovascular diseases, and immune system disorders. Research on the c-Kit gene can aid scientists in better understanding its functions in cell biology and pathophysiology, offering new insights for diagnosis and treatment of related diseases.
Storage	Store at 2-8°C short term (1-2 weeks).Store at ≤ -20°C long term. Avoid repeated freeze-thaw.
Formulation	Phosphate-buffered solution, pH 7.2.
Endotoxin	< 1 EU/mg, determined by LAL gel clotting assay

Data Examples

Flow cytometry

H_c-Kit(CD117) GNNK(+) CHO-K1 Cell Line (Catalog # GM-C36717) was stained with Anti-c-Kit(CD117) hlgG1 Antibody(briquilimab) (Catalog # GM-86543AB) or isotype control antibody, followed by anti-Human IgG APC-conjugated Secondary Antibody.




SampleID	Geometric Mean : FL11-H
CHO-K1 anti-CD117+APC-2nd Ab	1872
CHO-K1 H_c-Kit(CD117) GNNK(+) H_IgG+APC-2nd Ab	2038
CHO-K1 H_c-Kit(CD117) GNNK(+) anti-CD117+APC-2nd Ab	302330

Fig. FACS

Flow cytometry

Cynomolgus_c-Kit(CD117) GNNK(-) CHO-K1 Cell Line (Catalog # GM-C36581) was stained with Anti-c-Kit(CD117) hlgG1 Antibody(briquilimab) (Catalog # GM-86543AB) or isotype control antibody, followed by anti-Human IgG APC-conjugated Secondary Antibody.




SampleID	Geometric Mean : FL11-H
CHO-K1 anti-CD117+APC-2nd Ab	1729
CHO-K1 Cyno_c-Kit(CD117) GNNK(-) H_IgG+APC-2nd Ab	2220
CHO-K1 Cyno_c-Kit(CD117) GNNK(-) anti-CD117+APC-2nd Ab	4.54E5

Fig. FACS

Flow cytometry

H_c-Kit(CD117) GNNK(-) CHO-K1 Cell Line (Catalog # GM-C36580) was stained with Anti-c-Kit(CD117) hlgG1 Antibody(briquilimab) (Catalog # GM-86543AB) or isotype control antibody, followed by anti-Human IgG APC-conjugated Secondary Antibody.




SampleID	Geometric Mean : FL11-H
CHO-K1 anti-CD117+APC-2nd Ab	3847
CHO-K1 H_c-Kit(CD117) GNNK(-) H_IgG+APC-2nd Ab	2196
CHO-K1 H_c-Kit(CD117) GNNK(-) anti-CD117+APC-2nd Ab	264355

Fig. FACS

Flow cytometry

H_c-Kit(CD117) GNNK(-) HEK-293 Cell Line (Catalog # GM-C36579) was stained with Anti-c-Kit(CD117) hlgG1 Antibody(briquilimab) (Catalog # GM-86543AB) or isotype control antibody, followed by anti-Human IgG APC-conjugated Secondary Antibody.



SampleID	Geometric Mean : FL11-H
HEK-293 anti-CD117+APC-2nd Ab	6414
HEK-293 H_c-Kit(CD117) GNNK(-) H_IgG+APC-2nd Ab	2113
HEK-293 H_c-Kit(CD117) GNNK(-) anti-CD117+APC-2nd Ab	1.20E6

Fig. FACS

Bioactivity-ELISA

Cynomolgus c-Kit(CD117) Protein; His Tag (Catalog # GM-87120RP) was immobilized at 2 µg/ml (100 µL/well). Increasing concentrations of Anti-c-Kit(CD117) hIgG1 Antibody(briquilimab) (Catalog # GM-86543AB) were added.

Bioactivity-ELISA

0.2 µg Cynomolgus c-Kit(CD117) Protein; His Tag of per well




Fig. assay

Bioactivity-ELISA

Human c-Kit(CD117) Protein; His Tag (Catalog # GM-87117RP) was immobilized at 2 µg/ml (100 µL/well). Increasing concentrations of Human SCF Protein; mFc Tag (Catalog # GM-87653RP) were added.

Bioactivity-ELISA

0.2 µg Human c-Kit(CD117) Protein; His Tag of per well




Fig. assay