

Anti-IL17A hlgG1 Reference Antibody (Secubio)

Product Information

Product Name	Anti-IL17A hlgG1 Reference Antibody (Secubio)
Storage temp.	Store at 2-8°C short term (1-2 weeks).Store at $\leq -20^{\circ}\text{C}$ long term. Avoid repeated freeze-thaw.
Catalog# / Size	GM-87353MAB-1mg / 1 mg GM-87353MAB-5mg / 5 mg GM-87353MAB-25mg / 25 mg GM-87353MAB-50mg / 50 mg GM-87353MAB-100mg / 100 mg

Antibody Information


Expression System	CHO
Aggregation	< 5% as determined by SEC-HPLC
Purity	> 95% as determined by SDS-PAGE
Endotoxin	< 1 EU/mg, determined by LAL gel clotting assay
Sterility	0.2 μm Filtered
Target	IL17A
Clone	Secukinumab
Alternative Names	CTLA-8, CTLA8, IL-17, IL-17A, IL17, ILA17
Source/Isotype	Human IgG1(REEM), Kappa
Application	/
Description	This gene is a member of the IL-17 receptor family which includes five members (IL-17RA-E) and the encoded protein is a proinflammatory cytokine produced by activated T cells. IL-17A-mediated downstream pathways induce the production of inflammatory molecules, chemokines, antimicrobial peptides, and remodeling proteins. The encoded protein elicits crucial impacts on host defense, cell trafficking, immune modulation, and tissue repair, with a key role in the induction of innate immune defenses. This cytokine stimulates non-hematopoietic cells and promotes chemokine production thereby attracting myeloid cells to inflammatory sites. This cytokine also regulates the activities of NF-kappaB and mitogen-activated protein kinases and can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). IL-17A plays a pivotal role in various infectious diseases, inflammatory and autoimmune disorders, and cancer. High levels of this cytokine are associated with several chronic

inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.

Formulation phosphate-buffered solution, pH 7.4.


Data Examples

SDS-PAGE



On SDS-PAGE under reducing (R)/non-reducing(N-R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-HPLC



The purity of this product is more than 95% verified by SEC-HPLC